

LA-UR-21-28107

Approved for public release; distribution is unlimited.

Reconfiguring the metabolism of photosynthetic microbes for their development as biotechnological platforms Title:

Author(s): Gonzalez Esquer, Cesar Raul

Intended for: Seminar to Cemvita Factori Inc.

Issued: 2021-08-12



Seminar- Cemvita Factory Inc.

Reconfiguring the metabolism of photosynthetic microbes for their development as biotechnological platforms



Raul Gonzalez

8/13/2021



Utilization of the marine biodiversity to foster technological innovation




Bioluminescence (i.e. GFP)



Bioactive compounds (Limulus Amoebocyte Lysate)



Improved fisheries (i.e. Atlantic Salmon)







Bioremediation (i.e. *Alcanivorax*)


Algal Biofuels and Bioproducts

Cyanobacterial ultrastructure and its relation to genomic features (Pfams)




Photosynth Res (2016) 129:147– 157








Improve environmental stress tolerance

Increase tolerance to heat stress as a proxy for other stresses












Increase Product Output

Increase yields of an isoprene-evolving mutant via pathway engineering



Improve CO₂ fixation machinery Streamlined assembly of carboxysomes






Streamlined carboxysome


- Flexibility of subunits under environmental conditions
- Several subunits required for transfer to other organisms
- Complex assembly and regulation in other organisms
- Fixed stoichiometry of the composing domains
 - Increased portability (fewer number of proteins required)
- Simplified assembly and regulation (one core assembly protein) in other organisms
- Potential for improvement

Improve CO₂ fixation machinery

Streamlined assembly of carboxysomes



Research at LANL



3) Customized PHA production in cyanobacteria






Increasing carbon storage compounds in microalgae



Energy Efficiency & Renewable Energy

BIOENERGY TECHNOLOGIES OFFICE

Increasing carbon flux towards RuBisCO



Carbonic anhydrase overexpression:

- **CAH1** and **CAH6**: Shift the equilibrium towards bicarbonate (decrease CO₂ leakage).
- CAH3: Produce more CO₂ available for RuBisCO.
- CAH4: Mobilize bicarbonate pool from mitochondria to the CCM.
- BCA: Shown previously to increase photosynthetic efficiency in plants.

HCO₃- transporter overexpression:

• **HLA3 and LCIA:** Increase the bicarbonate flux towards the chloroplast.

Challenge overcome: Development of a transformation method for *Chlorella sorokiniana*




 Biolistic DNA delivery and antibiotic selection frequently yielded too many false positives.
 Screening practically impossible.

- ✓ Electroporation, Flow cytometry and Physiology
- Delivered fluorescent molecules by electroporation, screened by flow cytometry


Gate: (P1 in all)


Solution of the control of the c

Screened different culture conditions for recovery, selective pressure.



LANL team pushes boundaries of the state-of-technology






Overexpression mutants in CS:

- RG: CAH1, CAH4, HLA3, BCA, LCIA
- SN: SNRK
 Britton T. et al. (In preparation)
- EG: PiOxi, P5CS
- JM: Guar Gum synthesis
- CSM: Increasing Carbon sink


WSU: 2-Phenylethanol synthesis

Gene stacking likely required to increase carbon fixation rates





Single overexpression mutants generated in CS:

- RG: CAH1, CAH4, HLA3, BCA, LCIA
- No positive phenotype observed in single mutants. Likely, multiple gene overexpression required to cross all cellular barriers



Photosynthetic capacity studied by limiting carbon availability







discrimination tool developed in N. salina was implemented for measuring carbon use efficiency in Chlorella sorokiniana

Zidenga et al. Algal Research (2018)

Photosynthetic capacity studied by limiting carbon availability






developed in *N. salina* was implemented for measuring carbon use efficiency in *Chlorella sorokiniana*


Zidenga et al. Algal Research (2018)

... which we are using to evaluate Carbonic anhydrase (CA) mutants in Nannochloropsis gaditana (Matt Posewitz, CSM)


Headspace analysis of N. gaditana



Growth comparison of CA mutants vs WT



Presentation Outline



3) Customized PHA production in cyanobacteria





2) Increasing carbon storage compounds in microalgae




Energy Efficiency & Renewable Energy


BIOENERGY TECHNOLOGIES OFFICE

Multiscale Characterization: Increasing carbon storage molecules in algal biomass

 Algae accumulate carbon storage compounds (lipids, carbohydrates) after Nitrogen depletion



P. soloecismus



Multiscale Characterization: Increasing carbon storage molecules in algal biomass

 Algae accumulate carbon storage compounds (lipids, carbohydrates) after Nitrogen depletion

Carbohydrate

35

30

10

25

10

10

Protein


5

0


2 4 6 8 10 12 14 16 18 20 22 24

Time (days)

P. soloecismus



Proportion of lipids vs carbohydrates may vary under culture conditions



Carbohydrates (MBTH) Lipid (FAME)

40

30


20

0.35 ppt 3.5 ppt 17.5 ppt 35 ppt 52.5 ppt Salinity

Gonzalez-Esquer CR et al. Algal Research


Increasing Carbon Storage in Transgenic Algae


Overexpression of a H⁺-pumping pyrophosphatase (AVP1)




Avp1 in plants:

- PP_i homeostasis (metabolic regulation)
- Creates H⁺ gradient for ATP synthesis
- Phloem loading/sucrose transport
- Avp1 overexpression:
 - Increases biomass, salt tolerance, and drought tolerance in crops like cotton, tomato, rice, and wheat.
- Avp1 overexpression effect unknown in algae
- AVP1 overexpression increases carbohydrates under N deplete conditions










K Wright, R. Gonzalez, T. Dale

• AVP1 overexpression decreases intracellular pH and diminishes growth on low PO_4 conditions





 AVP1 overexpression improves biochemical composition (increased C storage) of algal biomass, likely due to PP_i synthesis (provoking an intracellular P-limiting condition).





1

BioManiac: Can we leverage biodiversity to match/increase performance of chemically developed materials?

We use **Polyhydroxyalkanoates** (PHA) as case study. PHAs are carbon storage molecules with can be used as bioplastic precursors.


PHA cyanobacteria ≠ PHA heterotrophic bacteria






Assessment of ML-Predicted Genes by Overexpression


 Gene targets were synthesized into Psba2KS vector and transformed into PCC 6803



PCR screening of genomic integration




Testing mutant genome




One band: segregated mutant; two bands: WT still present

Validation (CyanoGEBA-ML in PCC 6803)


Growth phenotypes in two different culture conditions





 PHB accumulation via BODIPY/Flow cytometry





Concluding remarks

- LANL is at the forefront of genetic engineering of microalgae
 - We have developed genetic engineering toolboxes for many strains, including Picochlorum soloecismus, Nannochloropsis salina, Chlorella sorokiniana
 - We have implemented such toolboxes for generating mutants with favorable phenotypes
- We have the opportunity to integrate metabolic features from different species (cyanobacteria>microalgae>plants) into a synthetic biology discovery and developmental platform
 - We can leverage the cyanobacterial metabolic "simplicity" to engineer complex organisms, i.e. for the production of renewable polymers
 - Unrivaled 'omics and machine learning scientific collaboration